

Further information

If you have any questions about your dental treatment please contact your local Haemophilia Centre for advice.
(Please see below).

Scottish Haemophilia Centres

Glasgow Comprehensive Care Haemophilia Centre

Glasgow Royal Infirmary
Castle Street
Glasgow
G4 0SF

Telephone: **0141 211 5127**

Dental Department: 0141 531 8916

Edinburgh Comprehensive Care Haemophilia & Thrombosis Centre

OPD1
Royal Infirmary of Edinburgh
Little France Crescent
Edinburgh EH16 4SA

Direct line: **0131 242 1270**

Out of hours contact: **0131 536 1000**

- ask to speak to the on-call haematologist.

Tayside Haemophilia Centre

Ward 34 Day Area
Ninewells Hospital & Medical School
Dundee DD1 9SY

Telephone: **01382 660111 bleep 4072**

Direct line / voicemail: **01382 632225**

**Dental team, based at Dundee Dental
Hospital 01382 381600.**

Grampian Area Haemophilia Centre Haematology Day Unit

Ward 307
Aberdeen Royal Infirmary
Foresterhill
Aberdeen, AB25 2ZN

Telephone: **0845 456 6000**

Special Care dentistry: **01224551901/551084**

Haemophilia Centre


Haematology Outpatients Department
Raigmore Hospital
Old Perth Road
Inverness, IV2 3UJ
Telephone: **01463 704020**



Review date: August 2017

mi • 279932

Guidance for Adult Dental Patients with Bleeding Disorders



You have been diagnosed as suffering from haemophilia or another bleeding disorder. It is very important that you receive regular dental care. This leaflet will be able to help you plan your treatment.

We advise everybody to see their dentist every six months if they have their own teeth or every two years if they are wearing dentures.

It is important to understand that you may not need to attend the hospital for all of your dental treatment.

Prevention of dental problems

Everybody is at risk of dental problems. However, the management of certain types of dental disease may cause bleeding. Therefore, preventing dental disease is very important. The most common diseases are periodontal disease (gum disease), dental decay and tooth wear.

Dental Treatment

The majority of your treatment will not cause significant bleeding and may be carried out by a general dental practitioner (dentist) or within the public dental service.

• Prevention Advice

At each appointment your dentist will talk to you about preventing dental problems. This may include information about your diet and advising on the use of fluoride toothpaste and other treatment where appropriate.

• Orthodontic Treatment

There are no problems with you receiving

orthodontic treatment using either a fixed or removable appliance. It is important that there are no sharp edges which might cause bleeding.

• Hygiene Therapy (e.g. scale and polish)

It is very unlikely that the routine cleaning of your teeth will cause any prolonged bleeding.

However, if you have serious inflammation of the gums (gingival) or gum disease, you may need factor replacement therapy to control any bleeding. You may need to be referred to the hospital for treatment.

• Removable Denture (e.g. false teeth)

Treatment is unlikely to cause any problems.

• Restorative Dentistry

Restorative dentistry, including the provision of crowns and bridges, can be carried out safely as long as your dentist follows the local anaesthetic guidelines.

• Root Fillings

The placement of root fillings should not cause any problems. On occasions it may be necessary to apply a medication into the root canal before completing the treatment.

• Extractions (having a tooth removed) and Surgery

In most cases this will have to take place in the hospital environment so that you can receive the appropriate factor replacement therapy and after care. Your haemophilia centre will confirm this.

Local Anaesthesia

If you need local anaesthetic, your dentist (general dental practitioner) will need to follow the local anaesthetic guidelines.

The following are local anaesthetic techniques and advice as to the need for factor replacement therapy.

Local Anaesthetic Techniques

Buccal infiltration - this is when you have anaesthetic near the side of the tooth next to the cheek. You should not need factor replacement therapy.

Intra-papillary injections - this is an injection into the gum between the teeth. You should not need factor replacement therapy.

Intraligamentary Injections - this is an injection into the space between the tooth and the bone. You should not need factor replacement therapy.

Lingual infiltration - this is when you have anaesthetic at the side of the tooth next to the tongue. **You may need factor replacement therapy.**

Inferior Dental Block - this is an injection where your bottom jaw including the lips and tongue are numb. **You may need factor replacement therapy.**

Please turn Over